

C-Reactive Protein as a Predictor of Active Coronary Disease and Worse Prognosis in Chest Pain Unit

Dan Oieru¹, Roy Beigel¹, Hanoch Hod^{1,2}, Pierre Chouraqui^{1,2}, Orly Goitein¹, Eli Konen¹, Roy Beinart¹, Paul Fefer¹, Ben-Ami Sela³, Ari Shamiss⁴, Jacob Or⁵, Shlomi Matetzky^{1,2}

¹Heart Institute, Sheba Medical Center, Tel Hashomer, Ramat Gan, ²Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, ³Biochemical Laboratory, ⁴Hospital Director, ⁵Emergency Department, Sheba Medical Center, Tel Hashomer, Ramat Gan, Israel

Objectives: In recent years chest pain units (CPU), have been proven to be an efficient modality for evaluating low-medium risk patients with chest pain. Although C-reactive protein (CRP) was proven to be an inflammation marker and an emerging cardiovascular and atherosclerosis risk factor, its prognostic value in CPU evaluation has yet to be determined.

Methods: The study comprised 621 consecutive patients admitted to the Tel Hashomer CPU. High sensitivity (hs)-CRP was determined in all patients within 8-12hrs of admission. As per CPU protocol all patients were monitored with an ST-analyzer monitor for ST changes and malignant arrhythmias. Patients were tested for troponin levels on admission and after 8-12hrs. Those ruled out for acute myocardial infarction (AMI) underwent non-invasive testing (stress SPECT thallium scan, stress echo, cardiac CT) for detection of significant coronary artery disease (CAD).

Results: 120 out of 621 patients (19%) had hs-CRP \geq 6 (high CRP); 501(81%) had CRP<6 (low CRP). There were no significant differences in baseline characteristics. 21% of high CRP patients were hospitalized compared with 11% of low CRP patients (p=0.006, OR 2.05, 95% CI 1.22-3.45). Although both high and low groups had similarly low incidents of AMI (2.5% vs. 2.2%, p=0.7), high CRP patients had a higher incidence of significant CAD by non-invasive testing (14% vs 7%, p=0.013) and eventually were more likely to undergo percutaneous coronary intervention for significant CAD.

Conclusion: CRP carries an important predictive value and may contribute to the fast tract chest pain evaluation of low-medium risk patients.

STEMI Culprit Lesion Angiographic Characteristics Affect Clinical Outcome

Atilla Bulut, Boris Varshitzky, Osman Aydin, Canbaba Arin, AbrahamTeddy Weiss,
Israel Gotsman, Chaim Danenberg, Chaim Lotan

Heart Institute, Hadassah University Hospital, Jerusalem, Israel

Background:

Culprit lesion characteristics may influence the technical approach and clinical outcome in patients undergoing primary PCI due to AMI.

Methods:

We evaluated the angiographic characteristics and MACE in 344 consecutive patients undergoing primary PCI between January 2005 and July 2007 in Hadassah University Hospital. Based on angiographic criteria patients were divided to into 2 groups: complicated (group A) and simple simple (group B) culprit lesions.

Results:

Average pain to needle time was 126 minutes. Culprit lesion was localized in the proximal and middle segments of the artery in 256 patients (74%). TIMI O - I flow on admission was detected in 328 patients and TIMI II-III flow in 16 patients. Single, double and triple vessel disease occurred in 40%, 35% and 25% of patients, respectively. Anterior wall MI comprised 46% of the cases. Six patients did not undergo PCI of which two underwent urgent CABG. Cardiogenic shock on admission occurred in 5% of the patients, 4% had malignant arrhythmia and 1% had mechanical complications due to AMI. Group A included 192 patients with large arteries (n=6), bifurcation lesions (n=44), large intracoronary thrombus (n=78), ostial lesions (n=9), acute on chronic occlusion (n=11), uncertain localizations (n=7), simultaneous occlusion of two arteries (n=5), acute instent thrombosis (n= 11), acute graft thrombosis (n=10) and need for multivessel PCI (n= 11). Group B included 152 patients with simple coronary occlusions. The differences in outcome are described in the following table:

	N	Angiographic success	Final TIMI flow II - III	Massive distal embolization	In hospital mortality
Group A	192	80%	80%	16%	2
Group B	152	98%	95%	7%	0
P		0.001	0.001	0.012	0.2

Conclusion:

In our single center experience, complicated lesions are the majority of lesions negotiated during primary PCI. These lesions are associated with lower angiographic success and worse clinical outcome.

Characteristics of Recurrent versus First AMI; Results from the ACSIS 2006 Survey

Leor Perl¹, Shlomo Behar³, Roseline Schwartz³, Morris Mosseri²

¹ Internal Medicine B, ² Cardiology Department, Meir General Hospital and the Sackler Faculty of Medicine, Kfar Saba, ³ The Neufeld Cardiac Research Institute, Sheba Medical Center, Tel Hashomer, Ramat Gan, Israel

Background: Little is known regarding the nature of a recurrent myocardial infarction. We sought to compare the characteristics of an event, in patients presenting with a recurrent MI (PMI – Previous MI group) to those with a first MI (FMI group).

Methods: Data was collected from the 2006 ACSIS (Acute Coronary Syndrome Israeli Survey) - a biannual survey on acute myocardial infarction performed in 26 intensive cardiac care units in Israel during a two-month period. 2074 patients suffered from MI, 30% of who were PMI patients

Results: Presented here are only results with $P < 0.05$.

Presentation: When compared to the FMI group, PMI patients had more risk factors and comorbidities. During the event less PMI patients presented with atypical angina (55 vs. 61%), more with heart failure (7 vs. 3%). Electrocardiographically, less PMI patients had ST elevation (26 vs. 51%); more patients had ST depression (27 vs. 18%), as well as lack of ECG changes (27 vs. 15%). **Treatment:** PMI patients were less likely to receive reperfusion therapy (medically or percutaneously); 12 vs. 25% of patients were treated with primary percutaneous coronary intervention (PCI); 47% compared to 69% had PCI during hospitalization, and 4% compared to 8% were treated with thrombolysis. **Outcomes:** PMI had more complications than FMI patients; heart failure (17 vs. 10%), pulmonary edema (13 vs. 7%), cardiogenic shock (6 vs. 4%), mitral regurgitation (5 vs. 2%) and acute renal failure (7 vs. 4%). Mortality was higher among patients with PMI; 4 vs. 2% at 7 days and 6 vs. 4% at 30 days.

Conclusions: 1. PMI patients were less likely to present with STEMI, possibly due to increased secondary preventive treatment for their disease, vessel collateral formation, and earlier self-referral. 2. Patients with PMI suffered more major complications during hospitalization than FMI patients, and experienced higher mortality rates.

Right Ventricular Dysfunction in Isolated Anterior ST Elevation Myocardial Infarction

Sergiy Bereza, Sergio Kobal, Harel Gilutz, Reuben Ilia, Doron Zahger

Cardiology, Soroka University Medical Centre, Faculty of Health Sciences, Ben Gurion University of the Negev, Beer Sheva, Israel

Background: Isolated occlusion of LAD does not usually affect right ventricular (RV) function. We describe for the first time a series of patients in who isolated anterior ST elevation MI (STEMI) was associated with right ventricular dysfunction.

Methods: We retrospectively identified all patients admitted to our medical center between 9/1996 and 1/2007 who had an anterior STEMI, had single vessel disease involving either the LAD, diagonal or ramus arteries who were found to have unexplained RV dysfunction by echocardiography within 24 hours of admission. Patients with any other cardiac pathology or pulmonary hypertension were excluded. Five patients fulfilled these criteria out of 438 patients who had a similar coronary anatomy but no RV dysfunction (1.14%). Demographic, clinical, echocardiographic and angiographic characteristics were studied. Echocardiographic and angiographic data were reviewed by a “blinded” investigator.

Results: A blinded investigator confirmed the diagnosis of RV hypokinesis in all subjects. All patients with RV dysfunction were men and their mean age was 52 (range: 40 to 60) years. All of them had no prior cardiac history. All patients had moderate – severe LV dysfunction. Echocardiographic follow-up was available for one patient 5 months after admission; RV dysfunction was completely resolved and LV dysfunction significantly improved. The coronary angiograms of all subjects were reviewed; absence of disease in the RCA was confirmed but we could not identify a coronary basis for RV dysfunction.

Conclusion: Isolated anterior STEMI not involving the RCA may rarely lead to significant RV dysfunction in relatively young subjects.. The mechanism for this previously undescribed observation is unclear. Complete resolution of RV dysfunction is probably common.

Therapeutic Hypothermia in Survivors of out of Hospital Cardiac Arrest is also Beneficial after Cardiac Asystole.

Ron Joseph Leor-Librach¹, Shirly Gmach Fluk¹, Boris Gluzman¹, Samer Kasem¹,
Guy Sheineman^{1,2}, Simcha Brandis², Ibrahim Jasmawie¹, Vered Nachliely¹, Lina Levit¹

¹ ICCU, *The Heart Institute*, ² *The Cath Lab, The Heart Institute, Laniado Hospital,*
Netanya, Israel

Out of hospital cardiac arrest is a condition with high mortality and devastating morbidity. Patients who are found in ventricular fibrillation generally have a more favorable outcome than patients found in asystole, which at some places are considered hopeless. This has led many physicians to a less aggressive approach in patients found in asystole. Results from recent clinical trials suggest a beneficial effect of induced mild hypothermia on the neurological outcome and mortality in patients with out of hospital cardiac arrest, and this therapeutic modality is now part of the guidelines. During the last five years we induced mild hypothermia in 102 survivors of out of hospital cardiac arrest. During their stay in the ICCU, 38% regained consciousness and 20% died. 52 patients were initially found in ventricular fibrillation, 35 patients in asystole and the other patients in a variety of other conditions, mainly asphyxia and electromechanical dissociation. Consciousness was regained by 50% of the ventricular fibrillation patients, 31% of the asystole patients and two of the 9 COPD patients. We conclude that cardiac asystole is not a hopeless condition, and induced mild hypothermia may be beneficial in some of these patients.

Spontaneous Reperfusion in ST-Elevation Myocardial Infarction is Associated with more Distal Coronary Lesions

David Leibowitz, Penko Gerganski, Johannes Nowatzky, A. Teddy Weiss, David Rott
Cardiology, Hadassah-Hebrew University Medical Center, Jerusalem, Israel

Background: Spontaneous reperfusion (SR) of the infarct-related artery may occur in patients with ST-segment elevation myocardial infarction (STEMI). Limited data are available on angiographic characteristics of these patients. The objective of this study was to examine if there are differences in the distance of the culprit lesion from the coronary ostium in patients with STEMI with and without SR.

Methods: Patients who presented with acute STEMI within 12 hours after pain onset and who underwent coronary angiography were entered into the study. Measurement of the distance from the coronary ostium to the culprit lesion was performed. SR was defined as significant relief of chest pain with an at least 50% resolution of ST segment elevation on follow-up electrocardiograph

Results: A total of 469 patients with STEMI were included in the study of whom 77 met criteria for SR and 392 who did not. A highly significant difference was seen in ostial to culprit lesion distance with the culprit lesion in the SR group being more distal than those in the non-SR group (45 ± 22 mm vs 39 ± 20 mm, p-value < 0.009).

Conclusions: In conclusion, our findings demonstrate that the location of the culprit lesion in STEMI patients who undergo SR is more distal in the involved artery than in STEMI patients without SR. Our findings suggest that the relatively lower thrombus burden and greater amount of vasoreactivity in these lesions may increase the chances of SR.

Preadmission on the Ambulance Tirofiban Infusion for STEMI Patients on the Way to Primary Percutaneous Intervention.

Menachem Nahir, Yevgeni Hazanov, Daoud Qarawani, Diab Ganem, Moein Abboud,
Yonathan Hasin

Cardiovascular Institute, Cardiology, Padeh Hospital, Tiberia, Israel

Aim: Determining the feasibility and safety of pre admission diagnosis of STEMI by a mobile intensive care unit (MICU) team using telemedicine, and early administration of the GP IIb/IIIa inhibitor (Tirofiban) on the way to immediate PCI

Method and results: During the time February 2006 up to November 2007 27 patients with suspected STEMI were evaluated in the ambulance, using HeartView™ and ECG trans-telephonic Recorder/Transmitter to the cardiac care unit. The diagnosis was confirmed by the cardiologic on call in 24 of them and a loading dose of Tirofiban was infused. The study group included 22 males (91.2%), mean age $59.9 \text{ y} \pm 13.5$, HTN in 47%, DM 20%, smokers 47%, anterior wall infarct in 45.8% and cardiogenic shock in 2 patients (8.3%).

The drug was given 27 ± 6 minutes pre arrival to the hospital, and 33 ± 19 minutes pre initiation of the angiography. an ST segment resolution pre angiography was noted in 20.4% and initial TIMI grade flow of 2 or 3 was demonstrated in 40% of the patients.

All patients had successful PCI to the culprit artery. The hospitalization course was uneventful for all patients, and no excess bleeding complications were noted. All patients were discharge home after 4.4 ± 1.6 days.

For analysis, a comparison was done between the study group to consecutive 145 other STEMI patients treated with primary PCI during the same time period in our center. No statistically significant difference was found for clinical, laboratory and outcome parameters.

Conclusion: based on this small pilot study, pre hospital diagnosis of STEMI using telemedicine equipment, and initiating GP IIb/IIIa antagonist seems to be feasible and safe, yet a larger study is needed for evaluation of beneficial effect.

Pseudonormal Electrocardiogram During Evolving Myocardial Infarction and Severity of the Infarct Related Artery (IRA) Stenosis

Oscar Kracoff, Shelly Shumilov Klipper, Abraham Caspi
Cardiology, ICCU, Kaplan Medical Center, Rehovot, Israel

Pseudonormalization of negative T waves is a well known indicator of myocardial ischemia. Pseudo-normalization of T waves during evolving ST elevation myocardial infarction (STEMI) and its correlation with the IRA has not been described. We studied the coronary angiographies of 34 pts. with STEMI which were divided into 2 groups according to their ECG evolution. Group A consisted of 18 pts whom during the course of STEMI evolved to a normal ECG pattern within 24 hours of admission. Group B consisted of 16 pts with T wave inversion within 24 hours. All pts of group A had T wave inversion at a later time or after PCI of the IRA. The table presents a summary of the relevant data.

		Group A	Group B	P value
Patients		18	16	
Male		13	13	
Mean age		57,6	58,7	NS
IRA		14LAD, 3Cx,1RCA	12LAD,3Cx,1RCA	
%stenosis	LAD	91.8(SD=10.6)	85.3 (SD=11.7)	P<0.07
	Cx	86.7% SD=5.8	68.3 SD=16.1	P<0.07
	RCA	91% SD=8.2	70.5 SD=21.7	NS

Results: There were no demographic differences between both groups. The severity of coronary stenosis was significantly higher in group A for the LAD, Cx and RCA, however for the RCA the severe stenosis was not statistically significant since there was only one pt. in each group. Thus, we conclude that this regressive pseudo normal pattern is a transitional ischemic stage probably due to incomplete revascularization of the IRA. These findings may explain cases of myocardial infarction with apparently normal ECG at presentation.

ST Deviation Pattern in Acute Myocardial Infarction is not Related to Lesion Location

Rott David, Penko Greganski, Johannes Nowatzky, A. Teddy Weiss, Tova Chajek-Shaul,
David Leibowitz

Medicine, Cardiology, Hadassah-Hebrew University Medical Center, Jerusalem, Israel

Background: Myocardial infarction may be classified as ST-elevation MI (STEMI) or non-ST elevation MI (NSTEMI). In STEMI the culprit artery is usually occluded, whereas in NSTEMI it is usually patent. The location of the ruptured plaque may influence the MI type. Herein we examine whether the distance from the coronary ostium to the culprit lesion is different in STEMI as compared to NSTEMI.

Methods: We reviewed our database and selected patients who presented with an acute MI and underwent coronary angiography within 7 days of admission. The analysis included 754 patients of whom 514 had STEMI and 240 had NSTEMI. QCA was performed and the distance from the ostium of the coronary artery to the site of thrombosis was measured.

Results: For both STEMI and NSTEMI patients the first 50mm (LAD and LCX) or the first 60mm (RCA) of the coronary artery contained 75% of the culprit lesions. There were no significant differences in median distances from the vessel ostium to the site of thrombosis between STEMI and NSTEMI patients

Conclusions: the distance from coronary ostium to culprit lesion is similar in STEMI and NSTEMI patients. Culprit lesion location does not appear to influence the development of STEMI as opposed to NSTEMI.

Seasonal Variation in Myocardial Infarction is Limited to Patients with ST-Elevations on Admission

David Leibowitz, David Planer, A Teddy Weiss, David Rott

Cardiology, Hadassah-Hebrew University Medical Center, Jerusalem, Israel

Background: Previous studies have demonstrated a seasonal variation in the incidence of acute myocardial infarction (AMI) with an increase in cases during the winter months. These studies have not assessed whether ST-elevation MI and non-ST-elevation MI exhibit similar changes. The object of this study was to compare the seasonal variation of STEMI as compared to NSTEMI.

Methods: All patients who presented with AMI and underwent coronary angiography within 7 days of admission were identified via the institutional database. STEMI diagnosis required admission ECG demonstrating ST elevation in at least two contiguous leads. All acute MI's not meeting criteria for STEMI were defined as NSTEMI. Patients were divided into monthly and seasonal groups based on the date of admission with MI.

Results: A total of 784 patients were included; 549 patients with STEMI on the basis of the admission ECG and 235 with NSTEMI. When STEMI patients were analyzed by season there were 170 patients (31%) in the winter months, a statistically significant difference of excess MI. ($p < 0.005$). When NSTEMI patients were analyzed there were 62 patients (26%) in the winter with no statistically significant difference in the seasonal variation.

Conclusions: Our findings suggest that the previously noted seasonal variation in the incidence of AMI is limited to patients presenting with STEMI. This suggests there are important physiological differences between STEMI and NSTEMI, the nature of which remain to be elucidated.