

A Pilot Study on Physical Conditions Related to Right Ventricular Infarction and Cardiogenic Shock in Acute Myocardial Infarction (AMI)-PCI Data

Elyahu Stoupel^{1,3}, Abid Assali^{1,3}, Igal Teplitzky^{1,3}, Evgeny Abranson², Peter Israelevich⁴,
Ran Kornowski^{1,3}

¹ *Division of Cardiology, Rabin Medical Center,* ² *Epidemiology & Informatics Unit, Rabin Medical Center, Petach Tikva,* ³ *Sackler Faculty of Medicine, Tel Aviv University,*

⁴ *Department of Geophysics & Cosmic Sciences, Tel Aviv University, Tel Aviv, Israel*

In 1170 primary PCI for AMI (2000-2006) 123 (10.5%) were described as right ventricular infarction (RVI) and 102 (8.72%) accompanied by cardiogenic shock (CSh) -factors negatively affecting AMI prognosis. The **aim** of this study was to check if these two conditions are related by some specific environmental physical activity markers.

Methods: daily physical activity parameters geomagnetic (GMA)-I-IV levels and cosmic ray activity –by neutron activity on the Earth's surface (imp/min) were obtained for each day in the study period (n=2557), days of AMI with culprit artery at LAD, RCA, CRX - and at days with AMI –right ventricular and with cardiogenic shock.

Results: patient characteristics-it was significantly more men in the RVI group (p=0,0005) as in the CSh and, inverse, for women (p=0,06). Patients in the CSh group were older than in RVI (p<0,0002). Woman >80y. old were more often in the CSh group. (p=0,05). On days of both RVI and CSh it was higher CRA (neutron) activity as at the whole study period (p=0,0003 for RVI; p=0,018 for CSh) and higher as in LAD, RCA AMI for RVI, (p=0.006 -LAD, 0,027-for RCA; p=0.014 for CRX) and CSh-AMI (p=0,009 for LAD; p=0,029 for RCA; 0.089 for CRX). RVI took place more often than CSh at two highest levels of GMA. (p=0,06).

Conclusion: AMI is accompanied by RVI and CSh on days with higher CRA compared with other groups of AMI; RVI is more often seen with concomitant high levels of GMA than AMI with CSh. CSh patients were older than RVI and woman in CSh prevailed.

It can be presumed that higher neutron (CRA) activity is predisposing to expanded myocardial damage in AMI.

Cardiac Telemedicine from the Prospective of the Patient and the Physician: an Update

Michal Golovner¹, Yigal Yanay¹, Nomi Malov¹, Mayera Elizur¹, Mira Tamari¹, Arie Roth^{1,2,3}

¹ SHL - Telemedicine, Israel, ² Cardiology, Sourasky Medical Center, ³ Sackler Faculty of Medicine, Tel-Aviv University, Tel Aviv, Israel

Telemedicine, a direct offshoot of cutting-edge technology, is assuming an important place in modern healthcare. 'SHL' telemedicine provides professional care to subscribers who contact its monitor center by telephone, describe their symptoms to the professional staff, and transmit their real-time ECGs (lately by cardiobeeperes with cellular telephone transmission). These data, combined with those stored in a central computer for instant access, determine individually appropriate therapeutic measures, including the dispatch of mobile intensive care units.

The telecardiological system monitors blood pressure, pulmonary peak flow, and selected cardiac parameters. The new 'Telemarker' device for transtelephonically measuring cardiac markers in whole blood was found safe and efficacious for independent use at the patient's home.

There are currently >60,000 'SHL' subscribers (age range 15-102 years). Since its establishment (1987), the service has handled >9,100,000 calls and ECG transmissions: solely telephonic medical assistance was required by >85% of the callers. SHL's CHF program led to 66% fewer hospitalizations.

Telemedicine services improved diagnosis and benefitted survivors of AMIs by decreasing mortality at one year from discharge following infarction. In addition numerous cases of unexplained episodes of weakness and pre-syncopacal episodes were also diagnosed as arising from rhythm and conduction disturbances.

Transtelephonic instruction to laymen who provided basic life support measures until professionals arrived at the scene resulted in a 10% survival rate of otherwise terminal patients.

Accurate diagnosis in the pre-hospital setting of patients with cardiac symptoms enhances identification of those with acute coronary syndromes, conserves resources by appropriate triage, and improves patients' motivation to seek medical assistance.

The Value of Computed Tomography Coronary Angiography before Invasive Coronary Angiography in Patients after Coronary Artery Bypass Grafts

Ghassan Zaid¹, Nabeel Makhoul¹, Arie Shefer¹, Amin Hasan¹, Samuel Rouchfleish¹,
Zeina Abdel-Rauf², Uri Rosenschein¹

¹ Cardiology Department, Bnai Zion Medical Center, Technion Institute Faculty of Medicine, ² Department of Radiology, Bnai Zion Medical Center, Haifa, Israel

Background: Coronary imaging using multidetector computed tomography (MDCT) might be an initial non-invasive modality for the diagnosis of coronary artery disease. In patients after coronary artery bypass grafts (CABG) the value of MDCT before invasive coronary angiography (ICA) had not been tested.

Objective: To examine the value of MDCT in patients after CABG who are planned for ICA with optional percutaneous coronary intervention (PCI).

Methods: We enrolled patients post CABG, high risk for ICA, who were admitted to our department for ICA due to acute coronary syndrome or electively for angina and objective ischemia on stress testing. All patients before ICA had undergone 64-slice MDCT for the detection of culprit vessel stenosis. Patients who had MDCT finding of significant stenosis (diameter stenosis >50%) had undergone ICA with optional PCI.

Result: Seven patients, 5 male and 2 female, mean age 77 ± 5 years were enrolled till now. According to MDCT finding, 4 patients underwent successful PCI of target saphenous venous graft (SVG) (PCI group); 3 of them had selective imaging of target vessel only. Three patients in whom MDCT showed significant target vessel stenosis were, according to ICA, not suitable for PCI (non-significant stenosis in 2 and chronic total occlusion in 1 patient), (intention to treat group). One patient in this group had selective imaging of target vessel only. Mean fluorescence time in patients with selective imaging of target vessel was 7 ± 4 minutes (min) compared with 9.5 ± 12 min in those with non-selective imaging. The mean amount of contrast media that given during ICA was 90 ± 76 ml compared with 110 ± 45 in selective and non-selective imaging respectively. All patients early after PCI and during the follow-up were clinically stable and uneventful.

Conclusion: In our study patients, MDCT before ICA is feasible and safe. MDCT guided PCI was associated with high procedural success rate, less fluorescence time and less amount of contrast media. In 3 of 7 patients the MDCT were not accurate with ICA findings. For statistical power, our study is still in progress.

Mild Therapeutic Hypothermia Improves Outcome after Out-of-Hospital Cardiac Arrest

Jonathan Balkin, Tanya Weizmann, Adi Butnaru, Michael Naroditzki, Ivetta Sucholotzky, Roman Korenfeld, Dan Tzivoni

Cardiology Department, Intensive Coronary Care Unit, Shaare Zedek Medical Center, Jerusalem, Israel

Treatment of coma following resuscitation with mild hypothermia has been incorporated into the guidelines following publications showing improvement in survival and neurological status after resuscitation for ventricular fibrillation (VF). We present the results of this treatment in our coronary care unit from 02/2003- 10/2007.

Results: Thirty patients (80% male) between ages of 23-82yrs were treated with mild hypothermia (34°C) following resuscitation. 23 had VF, 1 had VT and 6 had asystole as initial rhythm. Twenty two patients (73%) survived to discharge from hospital, 16 (53%) were discharged in good condition or moderate neurological deficit, and 6 (20%) had severe anoxic brain damage or vegetative state.

Predictors for good outcome included age <65yrs, length of resuscitation, and number of DC shocks. Asystole was not a predictor of poor outcome as 5 of the 6 patients admitted with asystole were discharged home with good neurological function.

Conclusions: This series of patients treated with mild hypothermia confirms the results of the randomized trials and suggests that treatment with hypothermia improved the outcome of survivors of sudden cardiac death and its indication may be broadened to include patients admitted not only with VF but also after asystolic cardiac arrest.

Patients' Personality and Spouses' Ways of Giving Support: Which Contributes More to the Recovery after First ACS?

Noa Vilchinsky^{1,2}, Liat Haze-Filderman¹, Morton Leibowitz², Orna Reges², Nili Erez²,
Rivka Elal², Tamir Bental², Dafna Manor², Orna Levit², Abed Khaskia², Morris Mosseri²,
David Daniel²

¹ Psychology, Bar-Ilan University, Ramat Gan, ² Cardiology, Meir Medical Center, Kfar
Saba, Israel

Background: In married couples the support provided by the spouse can be an important element in the healing process of the patient. Few studies have looked at the spouses' styles for giving support in the clinical context of recovery from an acute coronary syndrome (ACS). Coyne and Smith (1991, 1994) categorized three possible support styles of the spouse: active engagement, protective buffering and overprotection. They predicted that only the active engagement would be beneficial for the patients, however this was not empirically verified. Based on the person-environment fit model (Martire, Druley, Stephens & Wojno, 2002), our assumption was that support from the spouse will be beneficial for the patient recovery only as a function of the patient own personality characteristics.

Objective: The current study's main goal was to explore, in a prospective format, the nature of the interaction, between the spouses' ways of giving support and the patients' attachment personality style, to recovery outcomes of patients six months after their first ACS.

Method: The target population consisted of Jewish men, with the diagnosis of first ACS, admitted to the cardiac care unit at Meir medical center, Kfar Saba, Israel, between March 2005 and July 2006 (N=216). The study sample consisted of all patients and spouses who agreed to participate in the study and who completed the study questionnaires at Time 1-during hospitalization (N= 77), and at Time 2- six months after hospitalization (N=69). At time 1, the spouses completed the *Ways of Giving Support Questionnaire* (WOGS; Bunnk, et al, 2006), and the patients completed the *Experiences in Close Relationships* scale (ECR; Brennan, Clark & Shaver, 1998). Outcomes measured were: depression and anxiety reduction, BMI reduction, blood lipids improvement, Hs-CRP reduction, smoking cessation, rehabilitation program attendance and resuming work

Results: No main effect was found for attachment style in the prediction of the dependent variables. Two main effects were found for ways of giving support: active engagement predicted decrease in patients' BMI and over-protectiveness predicted non attendance in rehabilitation programs. Ten significant interactions were found between the patients' attachment styles (anxiety or avoidance) and the three ways of giving support of the spouses (active engagement, protective buffering and overprotection). These interactions correlated mostly with the patients' behaviors (i.e. smoking cessation, rehabilitation attendance and resuming work).

Conclusions: The interpersonal dynamics determining appropriate behavior following a medical crisis are complex and are an important element for further study. It is recommended that cardiac rehabilitation programs take in to consideration the contribution of the spouses' supportive style to the behavior and recovery of the male patient.

Factors Affecting non Attendance in Cardiovascular Rehabilitation Programs.

Rubin Lable², Harel Gilutz¹, Benny Pinkas³, Zahger Doron¹

¹ Cardiology department, Soroka University Medical Center, Faculty of Health Sciences, Ben Gurion University of the Negev, ² Department of Health Systems Management, Ben Gurion University of the Negev, ³ Cardionegev, Beer Sheva, Israel

Background: Attendance in phase-II cardiac rehabilitation program (CPR) is poor, and dropout rates of patients who start participation is unacceptable.

Aim: To identify factors influencing patients' decisions to attend cardiac rehabilitation programs.

Methods: Semi-structured interviews were conducted with 33 patients attending CPR, compared with 25 dropout patients that contacted the CPR.

Results: The age was of attending patients were 59.3±8 years, 73% were males.

Most of attending patients, 82% (n=29) could identify a reason for the cardiac event vs. only 36% (n=9) among non attending (p= 0.0002). Social environment in the rehabilitation center was identified as supporting further attendance among 68% of attending patients. Employment and distance from rehabilitation center was a reason for not participating in 90% of non attending patients. Surprisingly, health improvement was a minor factor influencing attendance decision in both groups. Information and encouragement delivered by the GP's were minimal and non influential in both groups. Factors not affecting attendance were: gender, employment, physical activity, spouse support and income.

Comments: Strategies should be developed for encouraging greater attendance by training GP's to identify and encourage eligible patients for CRP. Rehabilitation at out of work hours may enhance attendance. Distance is still a major obstacle of attendance, suggesting the importance of home-based rehabilitation.

Use of Attenuation Correction to Improve Accuracy of SPECT Myocardial Perfusion Imaging. Initial Experience at Hadassah Hospital

Elisha, James Ouzan¹, Martine Klein², Roland Chisin², Chaim Lotan¹, Moche Boscher²

¹ Heart Institute, ² Department of Nuclear Medicine, Hadassah Hebrew University Medical Center, Jerusalem, Israel

Objective: To assess prospectively the impact of attenuation correction (AC) on SPECT myocardial perfusion imaging (MPI) diagnostic accuracy.

Patients: Thirty two patients were referred to MPI: 11 patients for chest pain without known ischemic heart disease and 21 because of persistent chest pain after myocardial infarction and/or revascularization,

Methods: In addition to routine MPI, an x-ray tomography (GEMS –Hawkeye CT) was performed. Slice thickness was 1 cm. The CT Hounsfield units were transformed, for each pixel, into the corresponding attenuation coefficient for the used radioisotope (Tl-201 or Tc-99m). Ordered subsets expectation maximization (OSEM) algorithm was used for SPECT computer reconstruction to generate images with and without AC. The AC and non AC images were interpreted separately by 2 nuclear medicine physicians. A 6 to 12 months follow up period included close clinical follow up and, if necessary, coronary angiography. A consensus panel composed of cardiologists and NM physicians correlated the SPECT AC and non AC interpretations with the clinical and angiographic data.

Results: 1) Eight patients with normal AC MPI experienced no cardiac event during the follow up period. Seven of them showed 9 perfusion abnormalities on the non- AC data: in the infero-septal wall (5/7), apex (2/7) and anterior wall (2/7, male pts). 2) Eleven patients had inferior wall abnormality on the non-AC images and underwent angiography. Seven of them with abnormality seen also on the AC images had significant coronary disease in the RCA/Circumflex arteries. The other four with normal AC images had normal angiography of those arteries. 3) Two women with anterior wall perfusion abnormality on both AC and non AC images had a normal LAD coronary artery angiography. 4) In 10 patients with multiple perfusion abnormalities in territories other than the inferior wall, the correlation of angiography was better with the AC than with the non-AC images.

Conclusions: AC MPI may not perfectly correct for anterior wall defects, in particular for breast artifacts. However AC greatly improves the diagnostic accuracy of MPI SPECT and should be considered as becoming standard practice.

A Novel Noninvasive Digital Arterial Pulse Wave Analysis During Deep Breathing as an Indicator of Significant Coronary Artery Disease – A Two Center Study.

Amos Katz^{1,2}, David Brosh^{3,4}, Yosi Blaer¹, Abid Assali^{3,4}, Jamal Jafari^{1,2}, Ian Orlov¹,
Ran Kornowski^{3,4}

¹ Cardiology, Barzilai Medical Center, Ashkelon, ² Faculty of Health Sciences, Ben Gurion University, Beer Sheva, ³ Cardiac Catheterization Laboratories, Cardiology, Rabin Medical Center, Petach Tikva, ⁴ "Sackler" Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

Background: Previous studies have indicated that fluctuations in pulse wave amplitude (PWA) might be a predictor of coronary artery disease (CAD). Therefore analysis of the respiratory modulation spectral response (RMR) of the PWA during deep breathing at a frequency of 0.1 Hz can be performed by the novel Cardiometer® device, utilizing a photo plethysmograph sensor, attached to the patient's finger and proprietary software. We evaluated this noninvasive simple test as an indicator of significant CAD

Methods: The RMR results of 195 consecutive pts; (mean age 63.1 ± 11.6 years, 73% male) referred for coronary angiography in two academic medical centers (97 pts in Rabin MC and 98 pts in Barzilai MC), were compared with their coronary angiography results. Patients with luminal stenosis of ≥ 70%, or left main stenosis ≥ 50% were classified as having significant CAD. The Cardiometer RMR test was performed in the recovery room of the catheterization laboratory prior to the procedure, excluding CABG and STEMI. RMR was analyzed after baseline 20 seconds spontaneous breathing, followed by 70 seconds of guided deep breathing at 0.1 Hz. The test was repeated post procedure in 152 pts following PCI or diagnostic catheterization.

Results: The RMR (normal ranges 78% [best] to 0% [worst]) was significantly lower in pts with significant CAD (n=118) vs. pts with non-significant CAD (n=77) (14.57±23.15 vs. 38.75±19.68, P<0.001). The improvement in post procedure RMR was significantly higher in pts who had undergone successful PCI (n=107) as compared to pts (n=88) who had only diagnostic catheterization only (27.51± 23.45 vs. 20.00 ± 25.91, P<0.05). RMR was lower in pts (29/195) with recent MI (14.35 vs. 25.83, P<0.05). By using a receiver operating characteristic analysis, we identified an RMR<29% (sensitivity 81.4%, specificity 76.6%; positive and negative predictive values: 84.2%, 72.8%, respectively) to be the optimal cut-off value for predicting significant CAD. The clinical characteristics and the RMR results were similar in the two centers.

Conclusions: The novel digital PWA analysis test during deep breathing using the Cardiometer device is a simple, non-invasive bedside or office based test to detect significant CAD and to monitor patients with CAD post PCI.

Association between HbA1c and Erectile Dysfunction in Non-Diabetic Men with and without Ischemic Heart Disease

Dan Justo¹, Yaron Arbel¹, Arie Steinvil¹, Michael Kinori¹, Noa Mashav¹, Itzhak Herz²,
Jacob George², Amir Halkin², Gad Keren², Ariel Finkelstein², Shmuel Banai²

¹ Internal Medicine D, ² Cardiology, Tel-Aviv Sourasky Medical Center, Tel Aviv, Israel

Introduction: HbA1c is biomarker of endothelial dysfunction in non-diabetics. Endothelial dysfunction is implicated the pathophysiology of erectile dysfunction (ED) as well as ischemic heart disease (IHD). Although HbA1c levels are associated with ED in diabetic men, the association between HbA1c levels, ED and IHD has never been studied in non-diabetic men.

Methods: A total of 119 men underwent coronary angiography and filled the Sexual Health Inventory for Males questionnaire (SHIM). Excluded were men with diabetes mellitus and men with chronic inflammatory process. SHIM scores 21 or less represented ED.

Results: Included were 70 men, mean ages 59.4±13.3 years. Mean SHIM scores were 16.8±8.1 and mean HbA1c levels were 5.6±0.6% for the whole cohort. HbA1c levels were associated with SHIM scores ($r=-0.34$; $p=0.004$) for the whole cohort. HbA1c levels were not higher among men with IHD ($n=52$) compared with men without IHD ($n=18$), and among men with ED ($n=47$) compared with men without ED ($n=23$).

Conclusions: HbA1c levels are associated with ED among non-diabetic men with and without IHD. This finding sheds a new light on the clinical use of HbA1c in non-diabetics, and in men with IHD in particular.

Clinical Pharmacology in the Coronary Cathlab: Incidence of Drug Related Problems in Patients Undergoing Coronary Angiography

Bruria Hirsch^{1,2}, Shlomit Goren¹, Amnon Hoffman², Chaim Lotan¹, Haim Danenberg¹

¹ Cardiology, Hadassah Hebrew University Medical Center, ² School of Pharmacy, Hebrew University, Jerusalem, Israel

Background:

Drug related problems (DRP's) are underreported and consequently are an underestimated cause of morbidity and mortality. Recent epidemiologic evidence estimates that DRPs are the fourth to the sixth leading cause of death in USA. Important risk factors for adverse drug events or reactions included polypharmacy, female sex, drugs with a narrow therapeutic range, renal failure and advanced age. Over 50% of DRP is estimated as preventable. Interventional cardiology deals with relatively sick patients that are under polypharmacy therapy. The objectives of this study were to evaluate the frequency and the predictors for medication errors and adverse drug events in this high-risk population.

Methods:

Consecutive patients undergoing coronary angiography were interviewed and their records surveyed by a clinical pharmacologist for DRP (defined as adverse drug events, drug interactions, medication errors, dosage problems, frequency problems, drug- illness interaction, drug- age interaction, lack of drug in therapy and partial therapy problems). Detected DRP were further evaluated by a clinical cardiologist familiar with the patient.

Results:

165 patients were studied, 108 DRPs were detected in 64 patients (40%). DRPs were detected in older patients (65 ± 10 vs. 59 ± 1 , $p < 0.01$) that were treated with more drugs (7.46 ± 2.7 vs. 6.6 ± 2.5). Patients with DRPs had a higher incidence of co-morbidities including diabetes mellitus and previous organ transplantation and were significantly under-treated with beta blockers, statins and aspirin.

Conclusions:

Drug related problems are very frequent in patients undergoing coronary interventions and may significantly affect morbidity in this already sick population. Meticulous screening for DRP may reduce their incidence and improve patient care and well being.