



Christian
Doppler
Laboratory
for
Cardiac and Thoracic
Diagnosis & Regeneration



Christian Doppler
Forschungsgesellschaft

Secretome of apoptotic peripheral blood cells (APOSEC) attenuates microvascular obstruction in acute myocardial infarction

- (I) Role of platelet aggregation**
- (II) Role of vasodilation**

HJ Ankersmit

DIVISION OF THORACIC SURGERY
CHRISTIAN DOPPLER LABORATORY
for the Diagnosis & Regeneration of Cardiac and Thoracic Diseases
MEDICAL UNIVERSITY OF VIENNA
www.meduniwien.ac.at/applied-immunology



Christian
Doppler
Laboratory
for
Cardiac and Thoracic
Diagnosis & Regeneration



Christian Doppler
Forschungsgesellschaft

The following relevant financial relationships exist related to this presentation:

Medical University of Vienna/APOSCIENCE has claimed financial interest

WO2010070105-A1

WO2007121495A2


WO2010000820A1

WO2010079086A1

WO2010000819A1

EP10194916.2


Ankersmit HJ: APOSCIENCE AG, shareholder , Vienna based Start Up



APOSEC Reperfused AMI Large Animal Model



Pigs (~30kg)



Occlusion of the LAD



Intravenous administration of
porcine APOSEC
40 minutes after occlusion







According to
AHA & ESC
Guidelines




Reperfusion after 90 minutes

APOSEC Reperfused AMI Large Animal Model



Infarction Size after 30 days



What is APOSEC – Lessons learned from secretome analysis

Good evidence for treating AMI with stem cell secretome („conditioned medium“)



Paracrine action accounts for marked protection
of ischemic heart by Akt-modified mesenchymal
stem cells

Nat Med. 2005 Apr;11(4):367-8.


Secretome of conditioned stem cells and peripheral blood cells contains comparable levels of cytokines/chemokines

Eur Heart J. 2008 Dec;29(23):2851-8.

APOSEC – Secretome of irradiated peripheral monuclear cells



APOSEC attenuates myocardial injury during AMI



Basic Res Cardiol. 2011 Nov;106(6):1283-97.


Basic Res Cardiol. 2011 Jun;106(4):645-55.

Eur J Clin Invest. 2009 Jun;39(6):445-56.

No-reflow – Role of microvascular obstruction (MVO)

Despite infarct vessel patency ... disordered microvascular function and inadequate myocardial tissue perfusion are often present

Circulation. May 1992;85(5):1699-1705.



Circulation 2008;117:3152-3156.




Microvasculature of a porcine heart
provided by M Gyöngyösi

Aim of the study – Experimental setting

Impact of APOSEC on MVO

Influence of APOSEC on microvasculature in a porcine AMI model






Finding the mode of action: Effects on platelets and vasodilation *in vitro*

in vitro experiments

- | | | | | |
|------|----------------------|-------|--------------|--------------|
| (I) | aggregometer | FACS | ELISA | western blot |
| (II) | coronary ring assays | ELISA | western blot | |








Results I – APOSEC reduces MVO during AMI

MVO = hypoenhanced areas in late-enhanced MRI images




triplet		VT	
control	APOSEC	control	APOSEC
0.7 ±8.5	0.2 ±0.2	5.7 ±3.4	2.4 ±1.9
4.8 ±3.3	3.4 ±2.0	3.2 ±1.9	3.4 ±2.4

Results II – APOSEC mediates anti-aggregatory effects *in vitro*








n=8

Results III – mode of action – VASP phosphorylation





Thromb Haemost. 2011 Aug 1;106(2):253-62.



n=8


Results IV – platelet inhibition *in vivo*



No-reflow – Role of microvascular obstruction (MVO)

Despite infarct vessel patency ... disordered microvascular function and inadequate myocardial tissue perfusion are often present

Circulation. May 1992;85(5):1699-1705.






Circulation 2008;117:3152-3156.




Microvasculature of a porcine heart
provided by M Gyöngyösi

Results V – APOSEC mediates vasodilation






n=6-7


Results VI – APOSEC mediates vasodilation (coronary ring assay)




<http://www.currentprotocols.com>



Results VII – APOSEC mediates vasodilation (HUVEC culture)



<http://www.sciencecellonline.com>



Medical University Vienna

Christian Doppler Laboratory for Cardiac and Thoracic Diagnosis and Regeneration

Hendrik Jan Ankersmit
Thomas Schweiger
Michael Lichtenauer
Matthias Zimmermann
Andreas Mitterbauer
Lisa Wutzlhofer

Department of Cardiology

Mariann Gyöngyösi
Christian Plass

Institute of Physiology

Ivo Wolf
Alice Assinger

Department of Surgery



Patrick Starlinger



Faculty of Animal Science, Kaposvár University

András Jakab
Ervin Berényi
Zsolt Petrási




Results IX – Treatment of APOSEC leads to increased functional parameters in the long time



n=7-9


Results III – APOSEC mediates anti-aggregatory effects *in vitro*

COX and NO inhibition





n=3

Excursus – Is the effect of APOSEC limited to „healthy“ donors?



n=6-7

Results V – mode of action – VASP phosphorylation



n=3

Excursus – Is the effect of APOSEC limited to „healthy“ donors?

