

Effect of right atrial pacing on left ventricular function

¹David S. Blondheim, MD, ²Mark Kazatsker, MD,
³Yuval Konstantino, MD, ¹Aya Asif, MD, ²Simcha R. Meisel, MD, MSc,
²Michael Shochat, MD, ²Aaron Frimerman, MD,
²Avraham Shotan, MD, ³Noah Cohen-Liel, MD

¹Noninvasive Cardiology Unit, ²Division of Cardiology, Hillel Yaffe
Medical Center, Hadera; ³Units of Noninvasive Cardiology and
Electrophysiology, Soroka Medical Center, Beer Sheva



The authors have no conflict of interests to declare

Aim: To study the effect of incremental heart rates (HR) on LV longitudinal strain and other parameters of LV function.

- **Background:** A decrease in left ventricular (LV) longitudinal strain in stress echocardiograms has been proposed as a tool to detect ischemia. However, LV strain may be affected similarly by physiological changes induced by stress such as increased HR.



- **Methods:** We studied patients who had dual chamber pacemakers (mostly for sick sinus syndrome) with intact A-V conduction but without known coronary disease.

Patients were paced from the right atrium in 10 bpm increments until target HR was attained or until occurrence of A-V block or atrial fibrillation.

An echo study was performed and blood pressure (BP) measured at each stage. Global longitudinal strain was determined semi-automatically by the speckle tracking method.

- **Results:** Of 13 patients studied, 11 had adequate echo quality throughout the study (age 73.0 ± 3.6 years, 8 women).
- **Mean HR:** increased from 70.8 ± 10.4 to 123.8 ± 12.6 bpm, (mean increase: 52.3 ± 12.4 bpm).
- **Mean BP:** did not change (mean $132/70$ to $124/77$ mmHg, $p = \text{NS}$).
- **LV inflow and outflow stroke volumes:** (estimated by mitral and aortic velocity-time-integrals) decreased. Inflow: 27.2 ± 7.1 to 16.0 ± 4.9 cm; outflow: 22.0 ± 5.2 to 10.2 ± 2.8 cm, $p < 0.0001$ for both).
- **Cardiac output:** did not decrease.
- **LV longitudinal strain:** decreased in all patients (from mean $-16.5 \pm 3.7\%$ to $-14.1 \pm 3.3\%$, $P < 0.0001$).
- In figure: % decrease of strain from baseline (blue) and from previous stage (red).

Percent decrease of strain from baseline (blue) and from previous stage (red).



Changes in longitudinal strain of individual patients.

Conclusions

- There was an inverse relation between the increase in HR and longitudinal strain.
- This was not a result of changes in BP or CO which were unchanged by right atrial pacing.
- A decrease in longitudinal strain during stress echo studies may result from the increase in HR per se; it does not necessarily indicate ischemia.
- Normal ranges for decreased longitudinal strain secondary to increased HR during stress echo studies must be defined in order to facilitate a possible separation between the normal effects of increased HR and myocardial ischemia.

