



CLINICAL OUTCOMES OF ST-ELEVATION MYOCARDIAL INFARCTION PATIENTS ACCORDING TO GENDER

Leor Perl, Hana Vaknin-assa, Avital Porter,
Tamir Bental, Eli Lev, Ran Kornowski, Abid
Assali

Cardiology, Rabin Medical Center, Israel

All authors declare they have no conflict of interests in presenting this work

Background

- ▣ Recent literature had shown a greater risk for adverse clinical outcomes following ST-segment elevation myocardial infarction (STEMI) events in women undergoing percutaneous coronary intervention (PCI).
- ▣ We aimed to assess the impact of gender on clinical results following STEMI.

Methods


- ▣ We used our single center registry of patients treated for STEMI using primary PCI between January 2001 and July 2012.
- ▣ Procedural and angiographic results and clinical outcomes up to 2 years were collected and adjudicated for major cardiac adverse events.
- ▣ The presentation and clinical outcome of 1,935 patients with STEMI primary PCI was analyzed and compared according to gender.

Results -Baseline Characteristics

Parameter	Men	Women	P value
<u>Demographics</u>			
Age	59±12	68±13	0.001
Age>65y	26%	63%	<0.001
Diabetes	23%	37%	0.001
HTN	48%	63%	0.01
Anemia	24%	28%	0.03
Renal Failure	10%	24%	<0.001
<u>Presentation</u>			
Ant. AMI	45%	51%	0.07
Killip>1	12%	16%	0.2
LVEF<40%	39%	48%	0.002
Multiple vessel	59%	55%	0.3
DES	16%	10%?	0.007
Success	96%	92%	0.002
GP2b3a	73%	54%	0.001
Total	1584 (71.9%)	351 (18.1%)	

- All cases combined through and the 18.3% year. higher 29%
- In a different class were years

Survival According to Gender (Kaplan-Meier Curve)



and
on, stent
rgery
s.
one
re also
0.001;
ine
Killip
in/m²
1 or 2

- ▣ In this prospective, real world registry-based study, we have shown worse outcomes among female patients undergoing primary PCI for STEMI. However, after correction for co-morbidities, gender was no longer an independent predictor of outcomes.